

Anaphylaxis Emergency Action Plan

Patient Name:			Age:
Allergies:			
Asthma Yes (high risk for sev	ere reaction)	□ No	
Additional health problems beside	es anaphylax	dis:	
Concurrent medications:			
MOUTH THROAT* SKIN GUT LUNG* HEART*	itching, : itching, t itching, l vomiting shortnes	stoms of Anaphylaxis swelling of lips and/or tongutightness/closure, hoarsene hives, redness, swelling g, diarrhea, cramps ss of breath, cough, wheeze lse, dizziness, passing out	ie ss
°Some	symptoms c	sent. Severity of symptoms an be life-threatening. ACT	FASTI
Emergency Action Steps - DO NOT HES Inject epinephrine in thigh using (check one):		ITATE TO GIVE EPINEPHRII Adrenaclick (0.15 mg)	NE!
		☐ Auvi-Q (0.15 mg)	☐ Auvi-Q (0.3 mg)
		EpiPen Jr (0.15 mg)	EpiPen (0.3 mg)
		Epinephrine Injection, USI	P Auto-injector- authorized gener
		☐Other (0.15 mg)	Other (0.3 mg)
Specify others:			
MPORTANT: ASTHMA INHALERS			PENDED ON IN ANAPHYLAXIS.
2. Call 911 or rescue squad (before		•	
B. Emergency contact #1: home			
Emergency contact #2: home		work	cell
Emergency contact #3: home	_ 	work	cell
mments:	-		
ctor's Signature/Date/Phone Numb	er		
ent's Signature (for individuals un	da 40	2/2	